

# Indian Institute of Science

## UE 204: Midsemester Test

**Date:** 22/3/19.

**Duration:** 9.00 a.m.–10.00 a.m.

**Maximum Marks:** 100

1. A disc of inner radius  $a$ , outer radius  $b$ , and unit width along the  $z$ -direction is fixed rigidly at the inner boundary  $r = a$  and is subjected to a moment  $M$  on the outer boundary  $r = b$  by the application of suitable tangential tractions which are independent of  $\theta$  (see Fig. 1). The radial displacement is given by

$$u_r = c_1 \cos \theta + c_2 \sin \theta + \frac{c_3}{r},$$

where  $c_1$ ,  $c_2$  and  $c_3$  are constants to be determined. Assuming  $u_z = 0$  and treating the problem as two-dimensional (i.e.,  $u_r = u_r(r, \theta)$  and  $u_\theta = u_\theta(r, \theta)$ ), find the displacement components  $(u_r, u_\theta)$  in terms of  $(r, \theta, \lambda, \mu, M, a, b)$ . Make reasonable assumptions and state them clearly. The governing equations are

$$0 = (\lambda + \mu) \frac{\partial(\text{tr } \epsilon)}{\partial r} + \mu \left[ \frac{\partial}{\partial r} \left( \frac{1}{r} \frac{\partial(r u_r)}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u_r}{\partial \theta^2} + -\frac{2}{r^2} \frac{\partial u_\theta}{\partial \theta} \right],$$

$$0 = (\lambda + \mu) \frac{1}{r} \frac{\partial(\text{tr } \epsilon)}{\partial \theta} + \mu \left[ \frac{\partial}{\partial r} \left( \frac{1}{r} \frac{\partial(r u_\theta)}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u_\theta}{\partial \theta^2} + +\frac{2}{r^2} \frac{\partial u_r}{\partial \theta} \right].$$

The strain-displacement and constitutive relations are

$$\epsilon_{rr} = \frac{\partial u_r}{\partial r}, \quad \epsilon_{r\theta} = \frac{1}{2} \left[ \frac{1}{r} \frac{\partial u_r}{\partial \theta} + r \frac{\partial}{\partial r} \left( \frac{u_\theta}{r} \right) \right],$$

$$\epsilon_{\theta\theta} = \frac{1}{r} \left( \frac{\partial u_\theta}{\partial \theta} + u_r \right), \quad \text{tr } \epsilon = \epsilon_{rr} + \epsilon_{\theta\theta},$$

$$\boldsymbol{\tau} = \lambda(\text{tr } \epsilon) \mathbf{I} + 2\mu \boldsymbol{\epsilon}.$$




Figure 1: Hollow disk fixed rigidly at the inner boundary  $r = a$ , and subjected to a moment  $M$  on the outer boundary  $r = b$ .